A NEW ARRANGEMENT IN THE SUBFAMILY CYMINAE

(HEMIPTERA, LYGAEIDAE)

By Harry G. Barber, Collaborator, Entomology Research Branch, U. S. Department of Agriculture, Washington, D. C.

Stål, 1874, separated the Cyminae into two divisions, Cymaria and Ischnorhynchinaria. Since that time systematists have followed Stål and included the *Ninus* group of species in the tribe Ischnorhynchini. As this group of species have very distinctive characteristics readily separable from the Ischnorhychini, it should be raised to equal tribal rank and called Ninini.

KEY TO TRIBES OF THE SUBFAMILY CYMINAE

- - Head more or less deflexed anteriorly nearly or quite as wide as basal margin of pronotum; eyes either projected or stylate, remote from anterior angles of pronotum; ocelli set nearly as far apart as each is removed from eye. Costal margins of corium strongly contracted towards base; clavus more or less expanded apically; commissure subequal to length of scutellum. Body pilose (Ninus)

List of the genera under the respective tribes:

Cymini: Cymus Hahn; Cymodema Spin.; Arphnus Stål; Cyrtohamphus Stål; Gonystus Stål; Karamania Korm; Ontiscus Stål; Sephora Kirk. for C. criniger White; Nesocymus Kirk. for C. calvus White; Neocymus Van D.; Pseudocymus Van D.

Ischnorhynchini: Kleidoccrys Steph.; Polychismc Kirk. for Imbrius Stål; Crompus Stål; Neocrompus China; Rhiobia Bergr. (= Domiduca Dist.); Rhiophila Bergr.; Pylorgus Stål; Mesostates Reut.; Hovania Horv.; Syzgitis Bergr. Ninini: Ninus Stål (= Ossipaga Dist.); Cymoninus Bred.; Neoninus Dist.; Ninomimus Lundb.; Nesomartis Kirk.